

ON THE TORSION ANOMALOUS CONJECTURE IN CM ABELIAN VARIETIES

S. CHECCOLI AND E. VIADA

ABSTRACT. The Torsion Anomalous Conjecture (TAC) states that a subvariety V of an abelian variety A has only finitely many maximal torsion anomalous subvarieties. In this work we prove, with an effective method, some cases of the TAC when the ambient variety A has CM, generalising our previous results in products of CM elliptic curves. When V is a curve, we give new results and we deduce some implications on the effective Mordell-Lang Conjecture.

1. INTRODUCTION

Let A be an abelian variety embedded in the projective space and let V be a proper subvariety of A . Assume that both A and V are defined over the algebraic numbers.

Definition 1.1. *The variety V is a translate, respectively a torsion variety, if it is a finite union of translates of proper algebraic subgroups by points, respectively by torsion points.*

V is transverse, respectively weak-transverse, in A if V is irreducible and V is not contained in any translate, respectively in any torsion subvariety of A .

It is a classical problem in diophantine geometry to investigate the relationship between the above geometrical definitions and the arithmetical properties of the variety V . In this direction, there are several celebrated theorems such as the so-called Manin-Mumford, Mordell-Lang and Bogomolov Conjectures.

Recently E. Bombieri, D. Masser and U. Zannier in [BMZ07] suggested a new approach to this kind of investigations, introducing in particular the notion of torsion anomalous intersections.

Definition 1.2. *An irreducible subvariety Y of V is V -torsion anomalous if*

- *Y is an irreducible component of $V \cap (B + \zeta)$, with $B + \zeta$ an irreducible torsion variety;*
- *the dimension of Y is larger than expected, i.e.*

$$\text{codim } Y < \text{codim } V + \text{codim } B.$$

The variety $B + \zeta$ is minimal for Y if, in addition, it has minimal dimension. The relative codimension of Y is the codimension of Y in such a minimal $B + \zeta$.

We say that Y is maximal if it is not contained in any V -torsion anomalous variety of strictly larger dimension.

In [BMZ07], the authors formulate several conjectures. We state here one natural variant.

Conjecture 1.3 (TAC, Torsion Anomalous Conjecture). *For any algebraic subvariety V of a (semi-)abelian variety, there are only finitely many maximal V -torsion anomalous varieties.*

The TAC is well known to have several important consequences. It implies, for instance, the Manin-Mumford and the Mordell-Lang Conjectures; it is also related to model theory by work of B. Zilber and to algebraic dynamics by recent work of J. H. Silverman and P. Morton. In addition, R. Pink generalised it to mixed Shimura varieties.

Only few cases of the TAC are known: Viada [Via08] proved it for curves in a product of elliptic curves; Maurin [Mau08] for curves in a torus; Bombieri, Masser and Zannier [BMZ07] for varieties of codimension 2 in a torus. Habegger [Hab08] gave related results, under some stronger assumptions on V .

In [CVV13], we prove an effective TAC for maximal V -torsion anomalous varieties of relative codimension one in a product of CM elliptic curves. Our bounds are explicit and uniform in their dependence on V . As an immediate corollary, we prove the TAC for varieties of codimension 2, obtaining an elliptic analogue of the toric result in [BMZ07]. In this work we generalise our results to CM abelian varieties. In [CVV13], we also point out interesting relations between this kind of theorems and other relevant conjectures, such as the Zilber-Pink Conjecture and the above-mentioned ones.

Let $A \subseteq \mathbb{P}^m$ be an abelian variety with CM defined over a number field k and let k_{tor} be the field of definition of the torsion points of A . Let A be isogenous to a product of simple abelian varieties of dimension at most g . For a point $x \in A$, we denote by $\hat{h}(x)$ its canonical Néron-Tate height. For a subvariety $V \subseteq A$, we denote by $h(V)$ its normalised height and by $k_{\text{tor}}(V)$ its field of definition over k_{tor} (see Section 2). By \ll we denote an inequality up to a multiplicative constant depending on A . Our main result is the following:

Theorem 1.4. *Let $V \subseteq A$ be a weak-transverse subvariety of codimension $> g$. Then there are only finitely many maximal V -torsion anomalous subvarieties Y of relative codimension 1.*

Effective version: *More precisely, if $B + \zeta$ is minimal for Y , then for any positive real η there exist constants depending only on A and η such that:*

(1) *if Y is not a translate then*

$$\begin{aligned} \deg B &\ll_{\eta} (h(V) + \deg V)^{\text{codim } B + \eta}, \\ h(Y) &\ll_{\eta} (h(V) + \deg V)^{\text{codim } B + \eta}, \\ \deg Y &\ll_{\eta} \deg V (h(V) + \deg V)^{\text{codim } B - 1 + \eta}; \end{aligned}$$

(2) *if Y is a point, then*

$$\begin{aligned} \deg B &\ll_{\eta} ((h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}])^{\text{codim } B + \eta}, \\ \hat{h}(Y) &\ll_{\eta} (h(V) + \deg V)^{\text{codim } B + \eta} [k_{\text{tor}}(V) : k_{\text{tor}}]^{\text{codim } B - 1 + \eta}, \\ [k_{\text{tor}}(Y) : k_{\text{tor}}] &\ll_{\eta} \deg V [k_{\text{tor}}(V) : k_{\text{tor}}]^{\text{codim } B + \eta} (h(V) + \deg V)^{\text{codim } B - 1 + \eta}; \end{aligned}$$

(3) if Y is a translate of positive dimension, then

$$\begin{aligned} \deg B &\ll_{\eta} ((h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}])^{\text{codim } B + \eta}, \\ h(Y) &\ll_{\eta} (h(V) + \deg V)^{\text{codim } B + \eta} [k_{\text{tor}}(V) : k_{\text{tor}}]^{\text{codim } B - 1 + \eta}, \\ \deg Y &\ll_{\eta} \deg V ((h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}])^{\text{codim } B - 1 + \eta}. \end{aligned}$$

In addition, the torsion points ζ belong to a finite set of cardinality effectively bounded in terms of $\deg V$, $\deg B$ and constants depending only on A .

This theorem can be reformulated in the context of several other well known conjectures, as explained in the introduction of [CVV13].

The proof of Theorem 1.4 is split in two parts, depending on whether Y is a translate or not: in Section 4 we prove part (1) and in Section 5 we prove parts (2) and (3).

The main ingredients (see Section 2.3) needed for the proof of Theorem 1.4 are: Zhang's inequality, the Arithmetic Bézout Theorem by P. Philippon, our sharp Bogomolov type bound proved in [CVV12], the Relative Lehmer by M. Carrizosa. As usual, the CM hypothesis is due to the use of a Lehmer bound, known only for CM varieties. This result is only needed when Y is a translate, while case (1) of Theorem 1.4 holds with the weaker assumption that A has a positive density of ordinary primes, as required to apply a Bogomolov type bound (see [Gal10], p. 779). In particular, our method could treat the case of general abelian varieties, if the Lehmer and Bogomolov type bounds were known in such generality.

In Theorem 1.5, proved in Section 6, we expand our method in order to get some new effective results for curves in abelian varieties. This is particularly relevant, as bounds for the height in weak-transverse curves are hard to obtain. For instance, such bounds allow us to deduce some cases of the effective Mordell-Lang Conjecture, stated in Corollary 1.6. The two classical approaches to the effective Mordell-Lang Conjecture in abelian varieties are the Chabauty-Coleman and the Manin-Demjanenko methods. These methods require hypotheses that are similar to ours, but our result is of easier application and more explicit. Finally in Section 7, we give some generalisations to varieties in abelian varieties.

In particular we prove the following results. We fix an isogeny of the CM abelian variety A to the product $\prod_{i=1}^{\ell} A_i^{e_i}$ of non-isogenous simple factors A_i of dimension g_i . Since isogenies preserve finiteness, without loss of generality, we identify A with $\prod_{i=1}^{\ell} A_i^{e_i}$. If $H \subseteq A$ is a subgroup, then $H = H_1 \times \cdots \times H_{\ell}$, where $H_i \subseteq A_i^{e_i}$ is isogenous to $A_i^{f_i}$ for some $f_i \leq e_i$; therefore the matrix of the coefficients of the forms defining H has a structure of a block diagonal matrix, with entries in the endomorphism ring of the corresponding varieties. We can now state our effective result for weak-transverse curves, which is an example for the effective Zilber-Pink Conjecture.

Theorem 1.5. *Let $C \subseteq A = \prod_{i=1}^{\ell} A_i^{e_i}$ be a weak-transverse curve. Then the set*

$$\mathcal{S}(C) = C \cap \left(\bigcup_{H \in \mathcal{F}} H \right)$$

is a set of bounded Néron-Tate height, where \mathcal{F} is the family of all subgroups $H = \prod_{i=1}^{\ell} H_i \subseteq A$ such that

$$\text{codim } H_j > g_j \dim H_j$$

for at least one index j (here $\text{codim } H_j$ is the codimension of H_j in $A_j^{e_j}$).

More precisely, if $Y \in C \cap H$ then for any real $\eta > 0$ there exists a constant, depending only on A and η , such that

$$\hat{h}(Y) \ll_{\eta} (h(C) + \deg C)^{\frac{\text{codim } H_j}{\text{codim } H_j - g_j \dim H_j} + \eta} [k_{\text{tor}}(C) : k_{\text{tor}}]^{\frac{g_j \dim H_j}{\text{codim } H_j - g_j \dim H_j} + \eta}.$$

To prove Theorem 1.5 we first work in the projection on the j -th factor of A , and then we lift the construction to the variety A .

As an immediate consequence, we deduce the following corollary (proved in Section 6.1).

Let Γ be a subgroup of $A = \prod_{i=1}^{\ell} A_i^{e_i}$. Assume that the group $\overline{\Gamma}_i < A_i$ generated by the coordinates of the projections of Γ on the factors $A_i^{e_i}$ is an $\text{End}(A_i)$ module of rank t_i .

Corollary 1.6. *Let A be a CM abelian variety and let C be a weak-transverse curve in A . Let Γ be a subgroup as above, and suppose that $t_j < \frac{e_j}{g_j+1}$ for some index j . Then, for any positive η , there exists a constant depending only on A and η , such that the set $C \cap \Gamma$ has Néron-Tate height bounded as*

$$\hat{h}(C \cap \Gamma) \ll_{\eta} (h(C) + \deg C)^{\frac{e_j - t_j}{e_j - (g_j+1)t_j} + \eta} [k_{\text{tor}}(C) : k_{\text{tor}}]^{\frac{g_j t_j}{e_j - (g_j+1)t_j} + \eta}.$$

We remark that the corollary applies also to some Γ of infinite rank, indeed we only assume that the rank on one projection is small (see Remark 6.1).

2. PRELIMINARIES

2.1. Height and subgroups. We assume that all varieties are defined over the field of algebraic numbers.

Let A be an abelian variety with CM. We fix, up to an isogeny, a decomposition of $A = \prod_{i=1}^{\ell} A_i^{e_i}$ in simple factors, of dimension $\dim A_i = g_i$. We consider an embedding $i_{\mathcal{L}}$ of A in \mathbb{P}^m given by a symmetric ample line bundle \mathcal{L} on A . Heights and degrees corresponding to \mathcal{L} are computed via $i_{\mathcal{L}}$. More precisely, the degree of a subvariety of A is the degree of its image under $i_{\mathcal{L}}$; $\hat{h} = \hat{h}_{\mathcal{L}}$ is the \mathcal{L} -canonical Néron-Tate height of a point in A , and h is the normalized height of a subvariety of A as defined, for instance, in [Phi91]. Notice that if $x \in A$ is a point, then $\hat{h}(x) = h(x)$.

By Lemma 2.2 in [MW93], if A is an abelian variety defined over a number field k , then every abelian subvariety of A is defined over a finite extension of k of degree bounded by $3^{16(\dim A)^4}$; thus, without loss of generality, we assume that all abelian subvarieties of A are defined over k .

Let $B + \zeta$ be an irreducible torsion variety of A . Then $B = B_1 \times \dots \times B_{\ell}$ where $B_i \subseteq A_i^{e_i}$ is isogenous to $A_i^{f_i}$ for some integer $0 \leq f_i \leq e_i$.

There exists a natural correspondence between abelian subvarieties B of A , morphisms from A to $\prod_{i=1}^{\ell} A_i^{e_i - f_i}$, and matrices made of ℓ blocks, where the i -th block is an $(e_i - f_i) \times e_i$ -matrix with entries in the endomorphism ring of A_i . For details on such a correspondence see for instance [CVV12], Section 2.5. In short, the abelian subvariety B defines the projection morphism $\pi_B : A \rightarrow A/B$. The successive minima of B gives a matrix \mathcal{H}_B of the above type. By multiplication on the left, the matrix \mathcal{H}_B gives a morphism Φ_B from A to $\prod_{i=1}^{\ell} A_i^{e_i - f_i}$, with B the zero component of $\ker \Phi_B$.

By Minkowski's theorem, $\deg B$ is (up to constants depending only on A) the product of the squares of the norms of the rows of \mathcal{H}_B . In addition, B is the zero component of the zero set of the forms h_1, \dots, h_r corresponding to the rows of \mathcal{H}_B . We order the h_i 's by increasing degrees d_i , so that

$$d_1 \cdots d_r \ll \deg(B + \zeta) \ll d_1 \cdots d_r.$$

We also recall that, from [MW93], Lemma 1.3 and Lemma 1.4, if B is an abelian subvariety of A , and B^\perp is its orthogonal complement, then $\deg B^\perp \ll \deg B$, and therefore $\#(B \cap B^\perp) \ll (\deg B)^2$.

2.2. Torsion anomalous varieties. We recall some preliminary lemmas on torsion anomalous varieties used for our geometric constructions in the following sections.

Lemma 2.1 (Lemma 3.5 of [CVV13]). *Let Y be a maximal V -torsion anomalous variety and let $B + \zeta$ be minimal for Y . Then Y is weak-transverse in $B + \zeta$ (i.e. Y is not contained in any proper torsion subvariety of $B + \zeta$).*

Lemma 2.2 (Lemma 3.6 of [CVV13]). *Let Y be a maximal V -torsion anomalous variety, and let $B + \zeta$ be minimal for Y . Then Y is a component of $V \cap (B' + \zeta)$ for every algebraic subgroup $B' \supseteq B$, with $\text{codim } B' \geq \dim V - \dim Y$.*

The following lemma is due to Patrice Philippon [Phi12] and to certain properties of orthogonality in the Mordell-Weil groups studied by D. Bertrand [Ber86].

We recall that the *essential minimum* of a subvariety $X \subseteq A$ is defined as

$$\mu(X) = \sup\{\theta \in \mathbb{R} \mid \{x \in X(\overline{\mathbb{Q}}) \mid \hat{h}(x) \leq \theta\} \text{ is non dense in } X\}.$$

Lemma 2.3 (Philippon, Bertrand). *Let $H + Y_0$ be a weak-transverse translate in A , with Y_0 a point in the orthogonal complement H^\perp of H . Then $\mu(Y_0) = \mu(H + Y_0)$.*

We conclude with a remark on translations by torsion points.

Remark 2.4. *Notice that, for any subvariety X of A , translations by a torsion point ζ leave invariant the degree, the field of definition over k_{tor} and the normalised height of X (see also [Phi91], Proposition 9). In addition, if $Y \subseteq V \cap (B + \zeta)$ is V -torsion anomalous, then $Y - \zeta \subseteq (V - \zeta) \cap B$ is $(V - \zeta)$ -torsion anomalous. Therefore, without loss of generality, we can work in V or in $V - \zeta$, with the advantage, in this last case, that B is an abelian subvariety.*

2.3. Main ingredients. We recall here the main ingredients used in the proof of Theorem 1.4.

2.3.1. The Zhang Estimate. The following theorem follows from the crucial result in Zhang's proof of the Bogomolov Conjecture (see [Zha98]) and from the definition of normalized height.

Theorem 2.5 (Zhang). *Let $X \subseteq A$ be an irreducible subvariety.*

Then

$$\mu(X) \leq \frac{h(X)}{\deg X} \leq (1 + \dim X)\mu(X).$$

2.3.2. The Arithmetic Bézout Theorem. The following version of the Arithmetic Bézout Theorem is due to Philippon [Phi95].

Theorem 2.6 (Philippon). *Let X and Y be irreducible subvarieties of the projective space \mathbb{P}^n defined over $\overline{\mathbb{Q}}$; let Z_1, \dots, Z_g be the irreducible components of $X \cap Y$. Then*

$$\sum_{i=1}^g h(Z_i) \leq \deg X h(Y) + \deg Y h(X) + c(n) \deg X \deg Y,$$

where $c(n)$ is a constant depending only on n .

2.3.3. An effective Bogomolov Estimate. The following sharp Bogomolov bound is proved in [CVV12] and generalises a result of A. Galateau [Gal10].

Theorem 2.7 (Checcoli, Veneziano, Viada). *Let A be an abelian variety with a positive density of ordinary primes, and let Y be an irreducible subvariety of A transverse in a translate $B + p$. Then, for any $\eta > 0$, there exists a positive constant c_1 depending on A and η , such that*

$$\mu(Y) \geq c_1 \frac{(\deg B)^{\frac{1}{\dim B - \dim Y} - \eta}}{(\deg Y)^{\frac{1}{\dim B - \dim Y} + \eta}}.$$

2.3.4. A relative Lehmer Estimate. The following Lehmer bound is proved in [Car09].

Theorem 2.8 (Carrizosa). *Let A be an abelian variety with CM defined over a number field k , and let k_{tor} be the field of definition of all torsion points of A . Let P be a point of infinite order in A , and let $B + \zeta$ be the torsion variety of minimal dimension containing P , with B an abelian subvariety and ζ a torsion point. Then for every $\eta > 0$ there exists a positive constant c_2 depending on A and η , such that*

$$\hat{h}(P) \geq c_2 \frac{(\deg B)^{\frac{1}{\dim B} - \eta}}{[k_{\text{tor}}(P) : k_{\text{tor}}]^{\frac{1}{\dim B} + \eta}}.$$

3. FINITELY MANY MAXIMAL V -TORSION ANOMALOUS VARIETIES IN $V \cap (B + \text{Tor}_A)$

Let V be a weak-transverse variety in an abelian variety A . Let us fix an abelian subvariety B of A . In [CVV13], Lemma 3.9, we proved that there are only finitely many $\zeta \in \text{Tor}_{B^\perp}$ such that $V \cap (B + \zeta)$ has a maximal V -torsion anomalous component. In this section, Proposition 3.5, we prove that the number of such ζ is effectively bounded in terms of $\deg V$, $\deg B$ and some constants depending on A . We thank the referee for pointing out the effectivity question and for his useful comments.

The proof of such an effective result is based on an induction on the dimension of V , on Rémond's quantitative version of the Manin-Mumford Conjecture [Ré00] and on the effective bound for the degree of the maximal translates in a variety implied, for instance, by a result of Bombieri and Zannier [BZ96]. We first recall these results and some other well known bounds.

Recall that A is abelian variety and \mathcal{L} is a symmetric ample line bundle on A . We denote by $h_1(A)$ the projective height of the zero of A in the embedding associated with $\mathcal{L}^{\otimes 16}$ (as defined in [DP02], Notation 3.2) and by d_A the degree of the field of definition of A . If G is an abelian subvariety of A or a quotient of A , then $h_1(G)$ is bounded in terms of $h_1(A)$, $\deg A$, $\dim A$ and $\deg G$ (see [DP02], Proposition 3.9).

Moreover, in several works, Masser and Wüstholz, and then other authors, proved that for any abelian subvariety G of A , the degree of the field of definition of G is at most $3^{16(\dim A)^4} d_A$ (see [MW93] Lemma 2.2). Below, we sum up these bounds.

Estimate 3.1. *If G is an abelian subvariety of A or a quotient of A , then:*

- d_G is bounded in terms of d_A and $\dim A$;
- $h_1(G)$ is bounded in terms of $h_1(A)$, $\deg A$, $\dim A$ and $\deg G$.

For simplicity, in what follows we shall denote by $c(A) = c(\dim A, d_A, h_1(A), \deg A)$ any constant depending on $\dim A, d_A, h_1(A)$ and $\deg A$.

We recall the following consequence of Rémond's result, [Ré00] Theorem 1.2.

Estimate 3.2. *The number of irreducible components of the closure of the torsion of a weak-transverse variety V in an abelian variety A is effectively bounded as*

$$c(A)(\deg V)^{(\dim A)^{5(\dim V+1)^2}}.$$

Following the work of Rémond [Ré00] Theorem 2.1 and the results in [DP02] one sees that if G is an abelian subvariety of A or a quotient of A , then the corresponding constant $c(G)$ appearing in Estimate 3.2 is bounded only in terms of $\dim A, d_A, h_1(A)$, $\deg A$ and $\deg G$.

In our previous work joint with F. Veneziano [CVV13], Lemma 7.4, we gave an explicit version of a corollary of Lemma 2 in [BZ96]. This is a bound for the degree of the maximal translates contained in a variety, and so in particular for the degree of each component of the closure of the torsion. More precisely:

Estimate 3.3. *If V is weak-transverse in an abelian variety A , then the maximal translates contained in V have degree bounded by $c(A)(\deg V)^{2^{\dim V}}$.*

Notice that, if ζ is a torsion point such that $V \cap (B + \zeta)$ has a V -torsion anomalous component, then all the points in $\zeta + (B \cap B^\perp)$ share the same property. Indeed $B + \zeta = B + \zeta + (B \cap B^\perp)$. Clearly, we shall avoid such a redundancy and work up to points in $B \cap B^\perp$. Nevertheless, $|B \cap B^\perp| \ll (\deg B)^2$. This makes the following definition consistent.

Definition 3.4. *Let B be an abelian subvariety of an abelian variety A . Let V be a weak-transverse subvariety of A . We denote by $Z_{V,A}$ the set of torsion points $\zeta \in B^\perp / B \cap B^\perp$ such that $V \cap (B + \zeta)$ has a maximal V -torsion anomalous component Y_ζ .*

We point out that the set $Z_{V,A}$ also depends on B . However, in our proof B is fixed, while V and A vary. To simplify the notation we only indicate the dependence on V and A .

In the following proposition we estimate the number of points in $Z_{V,A}$. The number of maximal V -torsion anomalous components in $V \cap (B + \text{Tor}_A)$ is clearly estimated by $|Z_{V,A}| \deg V \deg B$.

Proposition 3.5. *Let B be an abelian subvariety of an abelian variety A . Let V be weak-transverse in A . Then the cardinality of $Z_{V,A}$ is effectively bounded in terms of $\deg V, \deg B$ and constants depending only on $\dim A, h_1(A), d_A$ and $\deg A$.*

Proof. Consider the projection

$$\pi_B : A \rightarrow A/B.$$

We recall that the degree of the image via π_B of a variety $X \subseteq A$ and the degree of the preimage via π_B of a variety $X \subseteq A/B$ only depend on $\deg X$, $\deg B$ and $\deg A$. In particular, $\deg \pi_B(V)$ is bounded in terms of $\deg V$, $\deg B$, $\deg A$ and $\deg A/B$ is bounded in terms of $\deg B$ and $\deg A$.

The proof of our proposition is done by induction on the dimension of V .

The base of our induction is the case of a curve, i.e. $\dim V = 1$. Then $\pi_B(V)$ is a weak-transverse curve in A/B , because V is weak-transverse in A . Moreover the points of $Z_{V,A}$ map to torsion points of $\pi_B(V)$. The number of torsion points of $\pi_B(V)$ is estimated using Estimate 3.2. Their preimage, which contains $Z_{V,A}$, has then cardinality effectively bounded in terms of $\deg V$, $\deg B$ and $c(A)$.

Suppose by inductive hypothesis, that the proposition holds for every variety V with $\dim V < n$. We then show that it holds for V of dimension n .

To prove our result, we are going to partition $Z_{V,A}$ into a finite union of subsets Z_X associated with irreducible subvarieties X of V of dimension $< n$. We then verify that such varieties X satisfy the assumption of the proposition; by the inductive hypothesis, we deduce that the cardinalities $|Z_X|$ are effectively bounded in terms of $\deg V$, $\deg B$ and $c(A)$.

Denote by $f : V \rightarrow A/B$ the restriction of π_B to V .

If f is dominant, then the generic fiber $F_p = V \cap (B + \tilde{p})$ has dimension

$$(1) \quad \dim F_p = \dim V - \operatorname{codim} B$$

here p belongs to an open subset of A/B and $f(\tilde{p}) = p$. The dimensional equation (1) shows that the generic fiber is not anomalous. Consider the subset V_π of A/B given by all points that do not have generic fiber. By the fiber dimension theorem (see for instance [Sha], Section 6.3, Theorem 7), this is a proper closed subset of $\pi_B(V)$ and its degree is effectively bounded in terms of $\deg V$, $\deg B$ and $c(A)$. Note that, the image of $Z_{V,A}$ via π_B is a subset of the torsion of V_π , indeed the fiber of a point in $\pi_B(Z_{V,A})$ is torsion anomalous and therefore does not satisfy the equality (1).

If f is not dominant, then set $V_\pi = \pi_B(V)$. Clearly, $Z_{V,A}$ is a subset of the torsion of V_π .

Note that in both cases:

(a) $\deg V_\pi$ is bounded in terms of $\deg V$, $\deg B$ and $c(A)$.

Let T_1, \dots, T_r be the isolated components of the closure of the torsion of V_π intersecting $\pi_B(Z_{V,A})$. Clearly

$$Z_{V,A} = \bigcup_{i=1}^r (\pi_B^{-1}(T_i) \cap Z_{V,A})$$

and

$$|Z_{V,A}| = \sum_{i=1}^r |(\pi_B^{-1}(T_i) \cap Z_{V,A})|.$$

From Estimate 3.2 and (a), the number r is effectively bounded in terms of $\deg V$, $\deg B$, $\deg A$ and $c(A)$. Thus we shall prove that, for every $1 \leq i \leq r$, the cardinality $|(\pi_B^{-1}(T_i) \cap Z_{V,A})|$ is effectively bounded in terms of $\deg V$, $\deg B$ and $c(A)$.

Let T be one of the above components. Define

$$W = \pi_B^{-1}(T) \cap V.$$

We have that:

- (i) $\deg W$ is bounded in terms of $\deg V$, $\deg B$ and $\deg A$. Indeed by Bézout's theorem, $\deg W \leq \deg \pi_B^{-1}(T) \deg V$. By Estimate 3.3, $\deg T$ is bounded in terms of the degree and the dimension of V_π and thus, by (a), in terms of $\deg B$, $\deg V$ and $c(A)$.
- (ii) $\dim W < n$, because V is weak-transverse in A and so it is not contained in $\pi_B^{-1}(T)$.
- (iii) For $\zeta \in \pi_B^{-1}(T) \cap Z_{V,A}$, each maximal V -torsion anomalous component Y_ζ of $V \cap (B + \zeta)$ is contained in W , indeed $\pi_B(Y_\zeta) = \pi_B(\zeta) \in T$.

By (iii), the variety W contains all the Y_ζ that we are counting, however W is not necessarily irreducible. Therefore we cannot hope to use the inductive hypothesis on W and we have to consider its irreducible components.

Let X_1, \dots, X_s be the irreducible components of W . For $\zeta \in \pi_B^{-1}(T) \cap Z_{V,A}$, we denote by Y_ζ any maximal V -torsion anomalous component of $V \cap (B + \zeta)$. By (iii), clearly each Y_ζ is contained in some X_i . We are going to count the number of Y_ζ contained in each X_i .

Denote by

$$Z_{X_j} = \{\zeta \in \pi_B^{-1}(T) \cap Z_{V,A} \mid X_j \text{ contains some } Y_\zeta\} / (B \cap B^\perp).$$

Then

$$\pi_B^{-1}(T) \cap Z_{V,A} = \bigcup_{j=1}^s Z_{X_j}.$$

The number s of irreducible components of W is bounded by $\deg W$. Thus, by (i), s is effectively bounded only in terms of $\deg V$, $\deg B$ and $c(A)$.

To conclude our proof we are left to bound in an effective way the cardinality of each Z_X , for X running over all irreducible components of W .

If X does not contain any Y_ζ , then $|Z_X| = 0$.

If $X = Y_{\zeta_0}$ for some $\zeta_0 \in Z_{V,A}$, then $|Z_X| = 1$.

Suppose that X strictly contains Y_{ζ_0} for some $\zeta_0 \in Z_{V,A}$. In this case we are going to show that $|Z_X| \leq |Z_{X - \zeta_0, \pi_B^{-1}(T) - \zeta_0}|$. Applying the inductive hypothesis, we estimate $|Z_{X - \zeta_0, \pi_B^{-1}(T) - \zeta_0}|$ in terms of $\deg V$, $\deg B$ and $c(A)$.

We first verify that $X - \zeta_0$ in $\pi_B^{-1}(T) - \zeta_0$ satisfies the assumption of the inductive hypothesis, that is the assumption of the proposition with $\dim X < n$. Observe that we need to translate by ζ_0 in order to obtain ambient varieties which are abelian varieties.

- The variety $\pi_B^{-1}(T) - \zeta_0$ is an abelian variety containing B . Indeed $B + \zeta_0$ is a subvariety of $\pi_B^{-1}(T)$ and $\zeta_0 \in \pi_B^{-1}(T)$.
- The variety $X - \zeta_0$ is weak-transverse in $\pi_B^{-1}(T) - \zeta_0$. Equivalently, by Remark 2.4, we show that X is weak-transverse in $\pi_B^{-1}(T)$. Since Y_{ζ_0} is a maximal V -torsion anomalous variety and X strictly contains Y_{ζ_0} , then X cannot be V -torsion anomalous. Recall that X is a component of $V \cap \pi_B^{-1}(T)$. Thus

$$(2) \quad \dim \pi_B^{-1}(T) - \dim X = \dim A - \dim V.$$

- If X were not weak-transverse in $\pi_B^{-1}(T)$, then $X \subseteq B_1 \cap V$ with $B_1 \subsetneq \pi_B^{-1}(T)$ a torsion variety. This contradicts relation (2).
- By (ii), $\dim X \leq \dim W < n$.

Thus, by inductive hypothesis, we get that:

$$|Z_{X-\zeta_0, \pi_B^{-1}(T)-\zeta_0}| \text{ is effectively bounded in terms of } \deg X, \deg B \text{ and } c(\pi_B^{-1}(T)).$$

We now show that by our construction $\deg X$ and $c(\pi_B^{-1}(T))$ only depend on $\deg V$, $\deg B$ and A .

- By (i), $\deg X \leq \deg W$ is effectively bounded in terms of $\deg V$, $\deg B$ and $c(A)$.
- By Estimate 3.2, we know that $\deg T$ is effectively bounded in terms of $\deg V_\pi$ and $\dim V$. Moreover, by (a), $\deg V_\pi$ is bounded in terms of $\deg V$, $\deg B$ and $c(A)$. Finally, Estimate 3.1 ensures that $d_{\pi_B^{-1}(T)}$ and $h_1(\pi_B^{-1}(T))$ are effectively bounded in terms of $\deg V$, $\deg B$ and $c(A)$.

Therefore

$$|Z_{X-\zeta_0, \pi_B^{-1}(T)-\zeta_0}| \text{ is effectively bounded in terms of } \deg V, \deg B \text{ and } c(A).$$

We finally prove that:

$$|Z_X| \leq |Z_{X-\zeta_0, \pi_B^{-1}(T)-\zeta_0}|.$$

We shall show that for every maximal V -torsion anomalous variety $Y_\zeta \subseteq X$, the variety $Y_\zeta - \zeta_0$ is a maximal $(X - \zeta_0)$ -torsion anomalous variety in $\pi^{-1}(T) - \zeta_0$.

Clearly $Y_\zeta - \zeta_0 \subseteq (X - \zeta_0) \cap (B + \zeta - \zeta_0)$. Since Y_ζ is V -torsion anomalous we have

$$\dim B - \dim Y_\zeta < \dim A - \dim V.$$

From this and (2) we obtain

$$\dim B - \dim Y_\zeta < \dim A - \dim V = \dim \pi_B^{-1}(T) - \dim X.$$

Thus $Y_\zeta - \zeta_0$ is a $(X - \zeta_0)$ -torsion anomalous variety.

In addition $Y_\zeta - \zeta_0$ is maximal: let $Y' \supset Y_\zeta - \zeta_0$ be a maximal $(X - \zeta_0)$ -torsion anomalous variety and let $B' + \zeta'$ be minimal for Y' . From (2), we have

$$\dim B' - \dim Y' < \dim \pi_B^{-1}(T) - \dim X = \dim A - \dim V.$$

Thus $Y' + \zeta_0 \subseteq V \cap (B' + \zeta' + \zeta_0)$ is V -torsion anomalous and containing Y_ζ . The maximality of Y_ζ as V -torsion anomalous implies $Y' + \zeta_0 = Y_\zeta$.

In conclusion, collecting all our bounds, we have proven that $|Z_{V,A}|$ is effectively bounded in terms of $\deg V$, $\deg B$ and $c(A)$. \square

4. NON TRANSLATE TORSION ANOMALOUS VARIETIES

Proof of Theorem 1.4 part (1). Let Y be a maximal V -torsion anomalous variety which is not a translate, and so of positive dimension. Let $B + \zeta$ be minimal for Y . We use the Arithmetic Bézout Theorem and the Bogomolov bound to prove that $\deg B$ is bounded only in terms of V and A , then we deduce the bounds for $h(Y)$ and $\deg Y$.

By Lemma 2.1 Y is weak-transverse in $B + \zeta$, and by assumption $\dim B = \dim Y + 1$; therefore Y is transverse in $B + \zeta$. Applying the Bogomolov estimate Theorem 2.7 to Y in $B + \zeta$ we get

$$(3) \quad \frac{(\deg B)^{1-\eta}}{(\deg Y)^{1+\eta}} \ll_{\eta} \mu(Y).$$

Let h_1, \dots, h_r be the forms of increasing degrees d_i such that $B + \zeta$ is a component of their zero set. We have that $r \leq \operatorname{codim} B \leq rg$ and

$$(4) \quad d_1 \cdots d_r \ll \deg(B + \zeta) = \deg B \ll d_1 \cdots d_r.$$

Consider the algebraic subgroup given by the first $h_1 \cdots h_{r-1}$ forms, and let B' be one of its irreducible components containing $B + \zeta$. Then by (4) we have

$$\deg B' \ll d_1 \cdots d_{r-1} \ll (\deg B)^{\frac{r-1}{r}}$$

and $\operatorname{codim} B' \geq \operatorname{codim} B - g$.

Since $\operatorname{codim} V \geq g + 1 = g + \dim B - \dim Y$, this implies that $\operatorname{codim} B' \geq \dim V - \dim Y$, and thus, by Lemma 2.2, Y is a component of $V \cap B'$.

We apply the Arithmetic Bézout Theorem to $V \cap B'$ and recall that $h(B') = 0$, because B' is a torsion variety; we get

$$(5) \quad h(Y) \ll (h(V) + \deg V) \deg B' \ll (h(V) + \deg V) (\deg B)^{\frac{r-1}{r}}.$$

Zhang's inequality, with (3) and (5), gives

$$\frac{(\deg B)^{1-\eta}}{(\deg Y)^{1+\eta}} \ll_{\eta} \mu(Y) \ll (h(V) + \deg V) \frac{(\deg B)^{\frac{r-1}{r}}}{\deg Y}.$$

Recall that Y is a component of $V \cap (B + \zeta)$. By Bézout's theorem $\deg Y \leq \deg B \deg V$, thus

$$(\deg B)^{1-\eta} \ll_{\eta} (h(V) + \deg V) (\deg B)^{\frac{r-1}{r}} (\deg B \deg V)^{\eta}$$

and therefore

$$(\deg B)^{\frac{1}{r}-2\eta} \ll_{\eta} (h(V) + \deg V) (\deg V)^{\eta}.$$

For η small enough we get

$$(6) \quad \deg B \ll_{\eta} (h(V) + \deg V)^{r+\eta} (\deg V)^{\eta};$$

this proves that the degree of B is bounded only in terms of V and A . Since the abelian subvarieties of bounded degree are finitely many, applying Proposition 3.5 we conclude that ζ belongs to a finite set of cardinality effectively bounded.

The bound on the height of Y is now given by (5) and (6)

$$h(Y) \ll_{\eta} (h(V) + \deg V)^{r+\eta} (\deg V)^{\eta}.$$

Finally, the bound on the degree is obtained from (6) and Bézout's theorem for the component Y of $V \cap B'$:

$$\deg Y \ll_{\eta} (h(V) + \deg V)^{r-1+\eta} (\deg V).$$

□

5. TORSION ANOMALOUS TRANSLATES

Proof of Theorem 1.4: parts (2) and (3). Let Y be a maximal V -torsion anomalous translate, with $B + \zeta$ minimal for Y .

We proceed to bound $\deg B$ and, in turn, the height and the degree of Y , using the Lehmer Estimate and the Arithmetic Bézout Theorem.

The variety $B + \zeta$ is a component of the torsion variety defined as the zero set of forms h_1, \dots, h_r of increasing degrees d_i , and

$$d_1 \cdots d_r \ll \deg B = \deg(B + \zeta) \ll d_1 \cdots d_r.$$

We have that $r \leq \text{codim } B \leq rg$.

Consider the torsion variety defined as the zero set of the first $r - 1$ forms h_1, \dots, h_{r-1} , and take a connected component B' containing $B + \zeta$, so that $\deg B' \ll d_1 \cdots d_{r-1} \ll (\deg B)^{\frac{r-1}{r}}$ and $\text{codim } B' \geq \text{codim } B - g$.

By Lemma 2.2, Y is a component of $V \cap B'$; indeed

$$\text{codim } B' \geq \text{codim } B - g = \dim A - g - \dim Y - 1 > \dim V - \dim Y - 1.$$

The proof is now divided in two cases, depending on $\dim Y$. If Y has dimension zero we use the Arithmetic Bézout Theorem and the Lehmer estimate; if $Y = H + Y_0$ is a translate of positive dimension, we can reduce to the zero dimensional case using some properties of the essential minimum.

Proof of part (2). Consider first the case of a maximal torsion anomalous point Y .

All conjugates of Y over $k_{\text{tor}}(V)$ are components of $V \cap (B + \zeta)$, they all have the same normalised height and their number is at least

$$[k_{\text{tor}}(V, Y) : k_{\text{tor}}(V)] \geq \frac{[k_{\text{tor}}(Y) : k_{\text{tor}}]}{[k_{\text{tor}}(V) : k_{\text{tor}}]}.$$

We then apply the Arithmetic Bézout Theorem in $V \cap B'$ obtaining

$$(7) \quad [k_{\text{tor}}(Y) : k_{\text{tor}}] \hat{h}(Y) \ll (h(V) + \deg V) [k_{\text{tor}}(V) : k_{\text{tor}}] (\deg B)^{\frac{r-1}{r}}.$$

Applying Theorem 2.8 to Y in $B + \zeta$, we obtain that for every positive real η

$$(8) \quad \hat{h}(Y) \gg_{\eta} \frac{(\deg B)^{1-\eta}}{[k_{\text{tor}}(Y) : k_{\text{tor}}]^{1+\eta}}.$$

Combining (8) and (7) we have

$$\begin{aligned} \frac{(\deg B)^{1-\eta}}{[k_{\text{tor}}(Y) : k_{\text{tor}}]^{\eta}} &\ll_{\eta} [k_{\text{tor}}(Y) : k_{\text{tor}}] \hat{h}(Y) \ll \\ &\ll (h(V) + \deg V) [k_{\text{tor}}(V) : k_{\text{tor}}] (\deg B)^{\frac{r-1}{r}}. \end{aligned}$$

For η small enough we obtain

$$(9) \quad \deg B \ll_{\eta} ((h(V) + \deg V) [k_{\text{tor}}(V) : k_{\text{tor}}])^{r+\eta} [k_{\text{tor}}(Y) : k_{\text{tor}}]^{\eta}.$$

Apply now Bézout's theorem to $V \cap B'$. All the conjugates of Y over $k_{\text{tor}}(V)$ are components of this intersection, so

$$(10) \quad \frac{[k_{\text{tor}}(Y) : k_{\text{tor}}]}{[k_{\text{tor}}(V) : k_{\text{tor}}]} \ll_{\eta} (\deg B)^{\frac{r-1}{r}} (\deg V).$$

Substituting (9) into (10) we have the last bound of part (2) in the statement.

Finally, we apply the Arithmetic Bézout Theorem to $V \cap B'$ to get

$$\hat{h}(Y) \ll (h(V) + \deg V)(\deg B)^{\frac{r-1}{r}} \ll_{\eta} (h(V) + \deg V)^{r+\eta} [k_{\text{tor}}(V) : k_{\text{tor}}]^{r-1+\eta}.$$

Having bounded $\deg B$, in view of Proposition 3.5 the points ζ belong to a finite set of cardinality effectively bounded.

Proof of part (3). Assume now that Y is a translate of positive dimension and write $Y = H + Y_0$, with H an abelian variety and Y_0 a point in H^{\perp} .

To bound $\deg B$ we can assume, without loss of generality, that $\zeta = 0$ (see Remark 2.4).

By Lemma 2.3,

$$(11) \quad \mu(Y_0) = \mu(H + Y_0).$$

Since the intersection $V \cap B'$ is defined over $k_{\text{tor}}(V)$, every conjugate of $H + Y_0$ over $k_{\text{tor}}(V)$ is a component of $V \cap B'$; as before, such components have the same normalised height and their number is at least

$$\frac{[k_{\text{tor}}(H + Y_0) : k_{\text{tor}}]}{[k_{\text{tor}}(V) : k_{\text{tor}}]}.$$

We apply the Arithmetic Bézout Theorem in $V \cap B'$ and we obtain

$$(12) \quad h(H + Y_0) \frac{[k_{\text{tor}}(H + Y_0) : k_{\text{tor}}]}{[k_{\text{tor}}(V) : k_{\text{tor}}]} \ll (h(V) + \deg V)(\deg B)^{\frac{r-1}{r}}.$$

By the Zhang's inequality, (11) and (12), we deduce

$$(13) \quad \mu(Y_0) \ll \frac{(h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}](\deg B)^{\frac{r-1}{r}}}{[k_{\text{tor}}(H + Y_0) : k_{\text{tor}}] \deg H}.$$

The lower bound for $\mu(Y_0)$ is derived as in the case of dimension zero.

Consider the smallest abelian subvariety H_0 of B containing Y_0 . Clearly H_0 is the irreducible component of $H^{\perp} \cap B$ containing Y_0 . Indeed, they are both one-dimensional abelian varieties containing the point Y_0 of infinite order.

By the definition of H_0 , $B = H + H_0$, and from [MW93], Lemma 1.2, we obtain

$$(14) \quad \#(H \cap H_0) \deg B \leq \deg H \deg H_0.$$

Moreover, from $H \cap H_0 \subseteq H \cap H^{\perp}$, we get

$$(15) \quad \#(H \cap H_0) \leq \#(H \cap H^{\perp}) \ll (\deg H)^2.$$

Applying Theorem 2.8 to Y_0 in H_0 we get that, for every positive real η

$$(16) \quad \mu(Y_0) = \hat{h}(Y_0) \gg_{\eta} \frac{(\deg H_0)^{1-\eta}}{[k_{\text{tor}}(Y_0) : k_{\text{tor}}]^{1+\eta}}.$$

We remark that

$$(17) \quad [k_{\text{tor}}(Y_0) : k_{\text{tor}}] \leq [k_{\text{tor}}(H + Y_0) : k_{\text{tor}}] \cdot \#(H \cap H_0)$$

because if $\sigma \in \text{Gal}(\overline{k_{\text{tor}}}/k_{\text{tor}}(H + Y_0))$ then $\sigma(Y_0) - Y_0 \in H \cap H_0$, so $[k_{\text{tor}}(Y_0) : k_{\text{tor}}(H + Y_0)] \leq \#(H \cap H_0)$.

Combining the upper bound and the lower bound for $\mu(Y_0)$ in (13) and (16), and using also (14), (15) and (17), for η sufficiently small, we have

$$(18) \quad \deg B \ll_{\eta} ((h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}])^{r+\eta}$$

where the dependence on $\deg H[k_{\text{tor}}(H + Y_0) : k_{\text{tor}}]$ has been removed applying Bézout's theorem to the intersection $V \cap B'$.

This also gives

$$\deg(H + Y_0) \ll_{\eta} (\deg V)((h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}])^{r-1+\eta}.$$

Finally, from (12), (18) and the trivial bound $[k_{\text{tor}}(H + Y_0) : k_{\text{tor}}] \geq 1$ we obtain

$$h(H + Y_0) \ll_{\eta} (h(V) + \deg V)^{r+\eta} [k_{\text{tor}}(V) : k_{\text{tor}}]^{r-1+\eta}.$$

Since we have bounded $\deg B$, we can conclude from Proposition 3.5 that the points ζ belong to a finite set of cardinality effectively bounded. \square

6. THE CASE OF A CURVE AND APPLICATIONS TO THE EFFECTIVE MORDELL-LANG CONJECTURE

Recall that $A = \prod_{i=1}^{\ell} A_i^{e_i}$, with A_i non-isogenous simple CM factors of dimension g_i . To prove Theorem 1.5 we essentially follow the proof of Theorem 1.4 part (2), working first in the projection on one factor, and then lifting the construction to the abelian variety A .

Proof of Theorem 1.5. Clearly all the points in $\mathcal{S}(C)$ are C -torsion anomalous; in addition, since C is a weak-transverse curve, each torsion anomalous point is maximal.

If $Y \in \mathcal{S}(C)$, then $Y \in C \cap H$, with $H = \prod_i H_i$ the subgroup containing Y which is minimal with respect to the inclusion.

Denote by Y_i the projection of Y on H_i and by C_i the projection of C on $A_i^{e_i}$. Let j be one of the index satisfying the hypothesis of the theorem. Assume first that Y_j is a torsion point and define $H' = A_1^{e_1} \times \cdots \times \{Y_j\} \times \cdots \times A_{\ell}^{e_{\ell}}$. Clearly $\deg H' \ll 1$ and $h(H') = 0$. Then, applying the Arithmetic Bézout Theorem to Y in $C \cap H'$, we get $\hat{h}(Y) \ll (h(C) + \deg C)$.

Assume now that Y_j is not a torsion point. Let $B_j + \zeta_j$ be a component of H_j containing Y_j . Clearly $\dim B_j = \dim H_j$ and $Y_j \in C_j \cap (B_j + \zeta_j)$ with $B_j + \zeta_j$ minimal for Y_j . Furthermore, Y_j is a component of $C_j \cap (B_j + \zeta_j)$ because C_j is weak-transverse and, by assumption, $\text{codim } H_j > g_j \dim H_j > 0$. This ensures that the matrix associated to $B_j + \zeta_j$ has at least 2 rows, which is necessary to apply the method.

We now sketch the proof which follows the proof of Theorem 1.4 part (2) and we give the relevant bounds.

The variety $B_j + \zeta_j$ is a component of the zero set of forms h_1, \dots, h_r of increasing degrees d_j with

$$d_1 \cdots d_r \ll \deg B_j = \deg(B_j + \zeta_j) \ll d_1 \cdots d_r,$$

and we have that $r = \text{codim } B_j / g_j = \text{codim } H_j / g_j$.

Consider the torsion variety defined as the zero set of h_1 , and let B'_j be one of its connected components containing $B_j + \zeta_j$; then $\deg B'_j \ll d_1 \ll (\deg B_j)^{\frac{1}{r}} = (\deg B_j)^{\frac{g_j}{\text{codim } B_j}}$.

From Theorem 2.8 applied to Y_j in $B_j + \zeta_j$, for every positive real η we get

$$(19) \quad \hat{h}(Y_j) \gg_{\eta} \frac{(\deg B_j)^{\frac{1}{\dim B_j} - \eta}}{[k_{\text{tor}}(Y_j) : k_{\text{tor}}]^{\frac{1}{\dim B_j} + \eta}}.$$

Notice that all conjugates of Y_j over $k_{\text{tor}}(C_j)$ are components of $C_j \cap B'_j$ and they all have the same height. Applying the Arithmetic Bézout Theorem to $C_j \cap B'_j$ and arguing as in the proof of Theorem 1.4, we have

$$(20) \quad \hat{h}(Y_j) \frac{[k_{\text{tor}}(Y_j) : k_{\text{tor}}]}{[k_{\text{tor}}(C_j) : k_{\text{tor}}]} \ll (h(C_j) + \deg C_j)(\deg B_j)^{\frac{g_j}{\text{codim } B_j}}.$$

Recall that $[k_{\text{tor}}(Y_j) : k_{\text{tor}}] \geq 1$. From (19) and (20) we get

$$(\deg B_j)^{\frac{\text{codim } B_j - g_j \dim B_j}{\text{codim } B_j \dim B_j} - \eta} \ll_{\eta} [k_{\text{tor}}(C_j) : k_{\text{tor}}](h(C_j) + \deg C_j)[k_{\text{tor}}(Y_j) : k_{\text{tor}}]^{\frac{1}{\dim B_j} - 1 + \eta}.$$

Since $\text{codim } B_j - g_j \dim B_j = \text{codim } H_j - g_j \dim H_j \geq 1$, for η sufficiently small this yields

$$(21) \quad \deg B_j \ll_{\eta} ([k_{\text{tor}}(C_j) : k_{\text{tor}}](h(C_j) + \deg C_j))^{\frac{\text{codim } H_j \dim H_j}{\text{codim } H_j - g_j \dim H_j} + \eta},$$

where we use $[k_{\text{tor}}(Y_j) : k_{\text{tor}}] \geq 1$ if $\dim B_j > 1$, and $[k_{\text{tor}}(Y_j) : k_{\text{tor}}] \leq [k_{\text{tor}}(C_j) : k_{\text{tor}}] \deg B \deg C_j$ if $\dim B_j = 1$.

We now lift the construction to A as follows. Define $H' = A_1^{e_1} \times \cdots \times B'_j \times \cdots \times A_{\ell}^{e_{\ell}}$. Clearly $\deg H' \leq \deg A \deg B'_j$ and Y is a component of $C \cap H'$. Applying the Arithmetic Bézout Theorem to $C \cap H'$ and using (21), we obtain

$$(22) \quad \hat{h}(Y) \ll (h(C) + \deg C) \deg H' \ll (h(C) + \deg C)(\deg B_j)^{\frac{g_j}{\text{codim } H_j}} \\ \ll_{\eta} (h(C) + \deg C)^{\frac{\text{codim } H_j}{\text{codim } H_j - g_j \dim H_j} + \eta} [k_{\text{tor}}(C) : k_{\text{tor}}]^{\frac{g_j \dim H_j}{\text{codim } H_j - g_j \dim H_j} + \eta}.$$

□

6.1. An application to the Effective Mordell-Lang Conjecture.

Proof of Corollary 1.6. Let $x \in C \cap \Gamma$. Let j be an index such that $\frac{e_j}{g_j + 1} > t_j$ and denote by (x_1, \dots, x_{e_j}) the projection of x in Γ_j .

Let $\gamma_1, \dots, \gamma_{t_j}$ be generators of the free part of $\bar{\Gamma}_j$. Then there exist elements $0 \neq a_k \in \text{End}(A_j)$ for $k = 1, \dots, e_j$, an $e_j \times t_j$ matrix M_j with coefficients in $\text{End}(A_j)$ and a torsion point $\zeta \in A_j^{e_j}$ such that

$$(a_1 x_1, \dots, a_{e_j} x_{e_j})^t = M_j(\gamma_1, \dots, \gamma_{t_j})^t + \zeta.$$

If the rank of M_j is zero, then (x_1, \dots, x_{e_j}) is a torsion point and so it has height zero.

If M_j has positive rank r_j , we can choose r_j equations of the system corresponding to r_j linearly independent rows of M_j . We use these equations to write the γ_k in terms of the x_k and we substitute these expressions in the remaining equations. We obtain a system of maximal rank with $e_j - r_j \geq e_j - t_j$ linearly independent equations in the variables x_1, \dots, x_{e_j} :

$$\begin{cases} m_{11}x_1 + \cdots + m_{1e_j}x_{e_j} = \xi_1 \\ \vdots \\ m_{e_j-r_j,1}x_1 + \cdots + m_{e_j-r_j,e_j}x_{e_j} = \xi_{e_j-r_j} \end{cases}$$

where $\xi_k \in A_j$ are torsion points and $m_{kl} \in \text{End}(A_j)$. These equations define a torsion variety $H_j \subseteq A_j^{e_j}$. Since $(g_j + 1)t_j < e_j$ we have $\text{codim } H_j > g_j \dim H_j$.

Then $x \in C \cap H$, where H satisfies the hypothesis of Theorem 1.5, which gives the bound for the height of x . □

Remark 6.1. Notice that it is possible to apply the corollary also to subgroups Γ whose rank is bounded only on one projection.

For example, let E_1, E_2 be two elliptic curves defined over \mathbb{Q} and such that $E_1(\mathbb{Q})$ is an abelian group of rank 1, and consider the product $A = E_1^4 \times E_2$.

Let C be a weak-transverse curve in A . Consider the subgroup $\Gamma = E_1(\mathbb{Q})^4 \times E_2(\overline{\mathbb{Q}})$ of A . Then Γ is not of finite rank, but, with the notation of the corollary, we have $g_1 = 1, e_1 = 4, t_1 = 1$ and $t_1 < \frac{e_1}{g_1+1} = \frac{4}{2}$.

The hypothesis of the corollary is therefore verified, and we have that

$$\hat{h}(C \cap \Gamma) \ll_{\eta} (h(C) + \deg C)^{3/2+\eta} [k_{\text{tor}}(C) : k_{\text{tor}}]^{1/2+\eta}.$$

7. FROM CURVES TO VARIETIES

We now adapt the proof strategy of Theorem 1.4 to obtain some new results for varieties V of dimension > 1 embedded in a power E^n of a CM elliptic curve. For subvarieties of general CM abelian varieties some technical condition arises. This makes a straightforward generalisation of our method of little interest.

For torsion anomalous varieties which are translates, the proof can be easily adapted, while for non translates a new argument is needed. Indeed, in this last case, the torsion anomalous variety is not transverse, but only weak-transverse in its minimal variety, a condition which is not sufficient to use the sharp Bogomolov bound.

The torsion varieties contained in V are already covered by the Manin-Mumford Conjecture, therefore we restrict ourselves to the V -torsion anomalous varieties which are not torsion.

Theorem 7.1. Let E be a CM elliptic curve defined over a number field k and let $n > 1$ be an integer. Denote by k_{tor} a field of definition of all torsion points of E .

Let $V \subseteq E^n$ be a weak-transverse variety. Let $Y \subseteq V \cap B + \zeta$ be a maximal V -torsion anomalous variety which is not a torsion variety, and let $B + \zeta$ be minimal for Y .

Set $b = \dim B$, $v = \dim V$ and $y = \dim Y$ and assume that $(n-b) > (v-y)(b-y)$. Then for any $\eta > 0$ there exist constants depending only on E^n and η such that:

(1) if Y is a point, then

$$\begin{aligned} \deg B &\ll_{\eta} ((h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}])^{\frac{(n-b)b}{(n-b)-vb} + \eta}, \\ \hat{h}(Y) &\ll_{\eta} (h(V) + \deg V)^{\frac{n-b}{(n-b)-vb} + \eta} [k_{\text{tor}}(V) : k_{\text{tor}}]^{\frac{vb}{(n-b)-vb} + \eta}, \\ [k_{\text{tor}}(Y) : k_{\text{tor}}] &\ll_{\eta} \deg V (h(V) + \deg V)^{\frac{vb}{(n-b)-vb} + \eta} [k_{\text{tor}}(V) : k_{\text{tor}}]^{\frac{n-b}{(n-b)-vb} + \eta}; \end{aligned}$$

(2) if Y is a translate of positive dimension, then

$$\begin{aligned} \deg B &\ll_{\eta} ((h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}])^{\frac{(n-b)(b-y)}{(n-b)-(v-y)(b-y)} + \eta}, \\ h(Y) &\ll_{\eta} (h(V) + \deg V)^{\frac{n-b}{(n-b)-(v-y)(b-y)} + \eta} [k_{\text{tor}}(V) : k_{\text{tor}}]^{\frac{(v-y)(b-y)}{(n-b)-(v-y)(b-y)} + \eta}, \\ \deg Y &\ll_{\eta} \deg V ((h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}])^{\frac{(v-y)(b-y)}{(n-b)-(v-y)(b-y)} + \eta}; \end{aligned}$$

(3) if Y is not a translate, then

$$\begin{aligned} \deg B &\ll_{\eta} ((h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}])^{\frac{(b-y)(n-b)}{(n-b)-(v-y)(b-y)} + \eta}, \\ h(Y) &\ll_{\eta} (h(V) + \deg V)^{\frac{(n-b)}{(n-b)-(v-y)(b-y)} + \eta} [k_{\text{tor}}(V) : k_{\text{tor}}]^{\frac{(v-y)(b-y)}{(n-b)-(v-y)(b-y)} + \eta}, \\ \deg Y &\ll_{\eta} \deg V ((h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}])^{\frac{(v-y)(b-y)}{(n-b)-(v-y)(b-y)} + \eta}. \end{aligned}$$

In addition the torsion points ζ belong to a finite set.

Proof of Theorem 7.1 part (1). Let Y be a maximal V -torsion anomalous point, with $B + \zeta$ minimal for Y .

We proceed to bound $\deg B$ and, in turn, the height of Y and the degree of its field of definition. To this aim we use the Lehmer bound in Theorem 2.8 and the Arithmetic Bézout Theorem.

Let $v = \dim V$ and $b = \dim B$. By Lemma 2.2, Y is a component of $V \cap B'$ where B' is, like in the proof of Theorem 1.4, the zero component of the torsion variety defined by the first v rows h_1, \dots, h_v of the matrix of B . Then $\text{codim } B' = v$ and $\deg B' \ll (\deg B)^{\frac{v}{n-b}}$.

We apply the Arithmetic Bézout Theorem to $V \cap B'$ to obtain

$$(23) \quad \hat{h}(Y) \ll \frac{(h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}]}{[k_{\text{tor}}(Y) : k_{\text{tor}}]} (\deg B)^{\frac{v}{n-b}}.$$

Applying the Lehmer estimate in Theorem 2.8 to Y in $B + \zeta$, instead, we have that for every positive real η

$$(24) \quad \hat{h}(Y) \gg_{\eta} \frac{(\deg B)^{\frac{1}{b} - \eta}}{[k_{\text{tor}}(Y) : k_{\text{tor}}]^{\frac{1}{b} + \eta}}.$$

From (23) and (24), and for η small enough, we get the bound for $\deg B$ if $b > 1$. If $b = 1$ we use Bézout's theorem to bound the factor $[k_{\text{tor}}(Y) : k_{\text{tor}}]^{\eta}$ as $((\deg B)(\deg V)[k_{\text{tor}}(V) : k_{\text{tor}}])^{\eta}$.

We then apply Bézout's theorem in $V \cap B'$ to bound $[k_{\text{tor}}(Y) : k_{\text{tor}}]$ and the Arithmetic Bézout Theorem in $V \cap B'$ to prove the bound for $\hat{h}(Y)$. Finally, from Proposition 3.5 it follows that the points ζ belong to a finite set of cardinality effectively bounded. \square

Proof of Theorem 7.1 part (2). Let $Y = H + Y_0$ be a maximal V -torsion anomalous translate of positive dimension, with minimal $B + \zeta$; assume also that $Y_0 \in H^{\perp}$.

We use the Lehmer bound and the Arithmetic Bézout Theorem to bound $\deg B$ and, in turn, the height and the degree of $H + Y_0$. In view of Remark 2.4, without loss of generality we can assume that $\zeta = 0$.

Let $b = \dim B$, $v = \dim V$ and $y = \dim Y = \dim H$. Clearly $v - y < n - b$ because Y is torsion anomalous.

As before, by Lemma 2.2 we have that Y is a component of $V \cap B'$, where B' is an irreducible torsion variety with $\text{codim } B' = v - y$ and $\deg B' \ll (\deg B)^{\frac{v-y}{n-b}}$.

Arguing as usual on the conjugates of $H + Y_0$ over $k_{\text{tor}}(V)$, we see that the intersection $V \cap B'$ has at least $[k_{\text{tor}}(H + Y_0) : k_{\text{tor}}]/[k_{\text{tor}}(V) : k_{\text{tor}}]$ components.

We apply the Arithmetic Bézout Theorem to the intersection $V \cap B'$, obtaining

$$(25) \quad h(H + Y_0) \ll (h(V) + \deg V)(\deg B)^{\frac{v-y}{n-b}} \frac{[k_{\text{tor}}(V) : k_{\text{tor}}]}{[k_{\text{tor}}(H + Y_0) : k_{\text{tor}}]}.$$

By Zhang's inequality, Lemma 2.3 and (25), we deduce

$$(26) \quad \mu(H + Y_0) = \mu(Y_0) \ll \frac{(h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}](\deg B)^{\frac{v-y}{n-b}}}{[k_{\text{tor}}(H + Y_0) : k_{\text{tor}}] \deg H}.$$

For the lower bound for $\mu(Y_0)$, the proof follows the case of $\dim Y = 0$. Let $H_0 = H^\perp \cap B$. By minimality of B we have that H_0 is a torsion variety of minimal dimension containing Y_0 , thus

$$\dim H_0 = \dim H^\perp + \dim B - n = (n - y) + b - n = b - y$$

As in Theorem 1.4 part (3), one can easily see that

$$(27) \quad [k_{\text{tor}}(Y_0) : k_{\text{tor}}] \leq [k_{\text{tor}}(H + Y_0) : k_{\text{tor}}] \cdot \#(H \cap H_0).$$

By the definition of H_0 , $B = H + H_0$ and from [MW93], Lemma 1.2, we get

$$(28) \quad \#(H \cap H_0) \deg B \leq \deg H \deg H_0.$$

In addition $H \cap H_0 \subseteq H \cap H^\perp$ thus

$$(29) \quad \#(H \cap H_0) \ll (\deg H)^2.$$

Applying Theorem 2.8 to Y_0 in H_0 we get that, for every positive real η

$$(30) \quad \mu(Y_0) = \hat{h}(Y_0) \gg_\eta \frac{(\deg H_0)^{\frac{1}{b-y}-\eta}}{[k_{\text{tor}}(Y_0) : k_{\text{tor}}]^{\frac{1}{b-y}+\eta}}.$$

Combining (26) and (30) we have

$$\frac{(\deg H_0)^{\frac{1}{b-y}-\eta}}{[k_{\text{tor}}(Y_0) : k_{\text{tor}}]^{\frac{1}{b-y}+\eta}} \ll_\eta \frac{(h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}](\deg B)^{\frac{v-y}{n-b}}}{[k_{\text{tor}}(H + Y_0) : k_{\text{tor}}] \deg H}$$

and hence, using (27), (28), (29) as it was done in Theorem 1.4 part (3), we get the bound for $\deg B$; notice that, if $b - y > 1$, the argument is in fact simpler, as we don't need to deal with the $[k_{\text{tor}}(Y_0) : k_{\text{tor}}]^\eta$ term.

Having obtained a bound for $\deg B$, the degree of $H + Y_0$ can be bounded applying Bézout's theorem to the intersection $V \cap B'$ and using $\deg B' \ll \deg B^{\frac{v-y}{n-b}}$. The bound for $h(H + Y_0)$, instead, is derived from (25) and the bound for $\deg B$. Finally, from Proposition 3.5 we conclude that the points ζ belong to a finite set of cardinality effectively bounded. \square

Proof of Theorem 7.1 part (3). Assume that $Y \subseteq V \cap (B + \zeta)$ is not a translate.

If Y is transverse in $B + \zeta$ the proof of Theorem 1.4 part (1) easily adapts to this case as well, yielding the desired bounds; let us then assume that Y is not transverse. Without loss of generality, we can assume $\zeta = 0$ (see Remark 2.4). Then Y is transverse in a translate $H_1 + Y_0 \subsetneq B$, with $Y_0 \in H_1^\perp$ and H_1 of minimal dimension.

We define $H_0 = B \cap H_1^\perp$, so that $B = H_1 + H_0$ and

$$(31) \quad \deg B = \deg(H_1 + H_0) \leq \frac{\deg H_1 \deg H_0}{\#(H_1 \cap H_0)}.$$

We set $y = \dim Y$, $v = \dim V$, $b = \dim B$, $h_1 = \dim H_1$ and $h_0 = \dim H_0 = b - h_1$.

Writing $Y = Y_1 + Y_0$ we have that $Y_1 \subseteq H_1$ is transverse in H_1 , because Y is transverse in $H_1 + Y_0$, and $Y_0 \subseteq H_0$ is transverse in H_0 , because B is minimal for Y .

By definition $Y_1 \subseteq H_1$ and $Y_0 \in H_1^\perp$. From Lemma 2.3 and the definition of essential minimum, we get

$$\mu(Y) = \mu(Y_1) + \hat{h}(Y_0).$$

As usual, the upper bound for $\mu(Y)$ is obtained using the Arithmetic Bézout Theorem in $V \cap B'$ for some abelian variety B' , constructed deleting $v - y$ suitable rows from B . All conjugates of Y are components of same height in $V \cap B'$. This gives

$$(32) \quad \mu(Y) \ll (h(V) + \deg V)(\deg B)^{\frac{v-y}{n-b}} \frac{[k_{\text{tor}}(V) : k_{\text{tor}}]}{\deg Y [k_{\text{tor}}(Y_1 + Y_0) : k_{\text{tor}}]}.$$

Moreover

$$(33) \quad [k_{\text{tor}}(Y_1 + Y_0) : k_{\text{tor}}] \#(H_1 \cap H_0) \geq [k_{\text{tor}}(Y_0) : k_{\text{tor}}]$$

because for every $\sigma \in \text{Gal}(\overline{k_{\text{tor}}}/k_{\text{tor}})$ which fixes $Y_1 + Y_0$, the difference $\sigma(Y_0) - Y_0$ lies in $H_1 \cap H_0$.

To obtain a lower bound for $\mu(Y)$ we either apply the Bogomolov bound to Y_1 in H_1 or the Lehmer estimate to Y_0 in H_0 . These give

$$(34) \quad \frac{(\deg H_1)^{\frac{1}{h_1-y}-\eta}}{(\deg Y)^{\frac{1}{h_1-y}+\eta}} \ll_\eta \mu(Y_1) \leq \mu(Y)$$

and

$$(35) \quad \frac{(\deg H_0)^{\frac{1}{h_0}-\eta}}{[k_{\text{tor}}(Y_0) : k_{\text{tor}}]^{\frac{1}{h_0}+\eta}} \ll_\eta \hat{h}(Y_0) \leq \mu(Y).$$

We now relate the left hand side to $\deg B$. Notice that one of the following relation holds: either

$$(i) \quad (\deg B)^{\frac{h_1-y}{b-y}} < \deg H_1$$

or

$$(ii) \quad (\deg B)^{\frac{h_0}{b-y}} \leq \frac{\deg H_0}{\#(H_1 \cap H_0)}.$$

Indeed if (i) and (ii) were both false, then

$$\deg B = (\deg B)^{\frac{h_1-y}{b-y} + \frac{h_0}{b-y}} > \frac{\deg H_1 \deg H_0}{\#(H_1 \cap H_0)},$$

which contradicts (31).

Assume that (i) holds. Then (32), (34) and (i) and the fact that $n - b > (v - y)(b - y)$ give the bound

$$\deg B \ll_\eta ((h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}])^{\frac{(b-y)(n-b)}{(n-b)-(v-y)(b-y)}+\eta},$$

where, if $h_1 - y = 1$, the factor $(\deg Y)^\eta$ has been removed applying Bézout's theorem to Y in $V \cap B$ and changing η .

Assume that (ii) holds. Then (32), (35) and (ii), the fact that $n - b > (v - y)(b - y)$ and (33) give the bound

$$\deg B \ll_\eta ((h(V) + \deg V)[k_{\text{tor}}(V) : k_{\text{tor}}])^{\frac{(b-y)(n-b)}{(n-b)-(v-y)(b-y)}+\eta}$$

where, if $h_0 = 1$ the dependence on $[k_{\text{tor}}(Y_0) : k_{\text{tor}}]$ can be removed using (33), bounding $[k_{\text{tor}}(Y) : k_{\text{tor}}]$ as $[k_{\text{tor}}(V) : k_{\text{tor}}] \deg V \deg B$ by the Bézout theorem

applied to Y in $V \cap B$ and observing that $\#(H_1 \cap H_0) \leq \#(H_1 \cap H_1^\perp) \ll (\deg H_1)^2 \ll (\deg Y)^{2h_1} \leq (\deg V \deg B)^{2h_1}$ because, since Y_1 is transverse in H_1 , $H_1 = Y_1 + \dots + Y_1$ (h_1 times), from which $\deg H_1 \ll (\deg Y)^{h_1}$.

So we have bounded $\deg B$. We obtain the bounds for $\deg Y$ and $h(Y)$ applying respectively the Bézout Theorem and the Arithmetic Bézout Theorem to the intersection $Y \subseteq V \cap B'$. Finally, Proposition 3.5 guarantees that the points ζ belong to a finite set of cardinality effectively bounded.

□

ACKNOWLEDGMENTS

We thank Francesco Veneziano for an accurate revision of an earlier version of this paper. We kindly thank the referee for his useful comments and corrections. Especially, we thank him for pointing out the effectivity question of Proposition 3.5.

REFERENCES

- [Ber86] Daniel Bertrand, *Relations d'orthogonalité sur les groupes de Mordell-Weil*, Séminaire de théorie des nombres, Paris 1984–85, Progr. Math., vol. 63, Birkhäuser Boston, Boston, MA (1986), pp. 33–39.
- [BMZ07] Enrico Bombieri, David Masser and Umberto Zannier, *Anomalous subvarieties—structure theorems and applications*, Int. Math. Res. Not. (2007), no. 19, pp. 1–33.
- [BZ96] Enrico Bombieri and Umberto Zannier, *Heights of algebraic points on subvarieties of abelian varieties*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **23** (1996), no. 4, pp. 779–792.
- [Car09] María Carrizosa, *Petits points et multiplication complexe*, Int. Math. Res. Not. (2009), no. 16, pp. 3016–3097.
- [CVV12] Sara Checcoli, Francesco Veneziano and Evelina Viada, *A sharp Bogomolov-type bound*, NYJM **18** (2012), pp. 891–910.
- [CVV13] ———, *On torsion anomalous intersections*. To appear in Rend. Lincei Mat. Appl. (2013).
- [DP02] Sinnou David and Patrice Philippon *Minorations des hauteurs normalisées des sous-variétés de variétés abéliennes II*, Comment. Math. Helv. **77** (2002), pp. 639–700.
- [Gal10] Aurélien Galateau, *Une minoration du minimum essentiel sur les variétés abéliennes*, Comment. Math. Helv. **85** (2010), no. 4, pp. 775–812.
- [Hab08] Philipp Habegger, *Intersecting subvarieties of \mathbf{G}_m^n with algebraic subgroups*, Math. Ann. **342** (2008), no. 2, pp. 449–466.
- [Mau08] Guillaume Maurin, *Courbes algébriques et équations multiplicatives*, Math. Ann. **341** (2008), no. 4, pp. 789–824.
- [MW93] David Masser and Gisbert Wüstholz, *Periods and minimal abelian subvarieties*, Ann. of Math. (2) **137** (1993), no. 2, pp. 407–458.
- [Phi91] Patrice Philippon, *Sur des hauteurs alternatives. I*, Math. Ann. **289** (1991), no. 2, pp. 255–283.
- [Phi95] ———, *Sur des hauteurs alternatives. III*, J. Math. Pures Appl. (9) **74** (1995), no. 4, pp. 345–365.
- [Phi12] ———, *Sur une question d'orthogonalité dans les puissances de courbes elliptiques*, Preprint (2012), <http://hal.archives-ouvertes.fr/hal-00801376>.
- [Rém00] Gaël Rémond, *Décompte dans une conjecture de Lang*, Invent. Math. **142** (2000), pp. 513–545.
- [Sha] Igor R. Shafarevich, *Basic Algebraic Geometry 1*, Springer-Verlag, 1972.
- [Via08] Evelina Viada, *The intersection of a curve with a union of translated codimension-two subgroups in a power of an elliptic curve*, Algebra Number Theory **2** (2008), no. 3, pp. 249–298.
- [Zha98] Shou-Wu Zhang, *Equidistribution of small points on abelian varieties*, Ann. of Math. (2) **147** (1998), no. 1, pp. 159–165.